



Apr 8th, 2:00 PM - 3:35 PM

Fisher Esterification of a Rhodium Substituted Keggin-Type Polytungstophosphate

Jeffrey W. Donelan
Illinois Wesleyan University

Rebecca Roesner, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Donelan, Jeffrey W. and Roesner, Faculty Advisor, Rebecca, "Fisher Esterification of a Rhodium Substituted Keggin-Type Polytungstophosphate" (2006). *John Wesley Powell Student Research Conference*. 32.

<https://digitalcommons.iwu.edu/jwprc/2006/posters2/32>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P18

**FISHER ESTERIFICATION OF A RHODIUM SUBSTITUTED KEGGIN-TYPE
POLYTUNGSTOPHOSPHATE**

Jeffrey W. Donelan and Rebecca Roesner*
Chemistry Department, Illinois Wesleyan University

Polyoxometalate (POM) chemistry is an emerging field with many applications in catalysis, electrochemistry, and medicine. Keggin-type POMs have the formula $[XM_{12}O_{40}]^n$, where M is an early transition metal and X can be almost any element. In recent years, chemists have become interested in attaching organic ligands to POMs for the purpose of creating immobilized polyoxoanion catalysts.¹ In our present work, a rhodium substituted Keggin-type POM with a pendent carboxylic functional group ($[PW_{11}O_{39}RhCH_2COOH]^{5-}$) was synthesized by previously reported methods.^{1,2} This POM was then converted to its methyl and ethyl ester derivatives through Fisher esterification with the appropriate alcohols. The ester products were then characterized by NMR and IR spectroscopies. A further effort is being made to immobilize the carboxylic acid functionalized polytungstophosphate through Fisher esterification with alcohol functionalized resin beads.

1. Wei, X. Dickman, M.H.; Pope, M.T. *J. Am. Chem. Soc.* **1998**, *120*, 10254-10255.
2. Unpublished results. Binder, J., Roesner, R. Illinois Wesleyan University.