



Apr 12th, 2:35 PM - 3:35 PM

Host-Guest Interactions Between Niobium Polyoxometalates and Neutral Aza-Macrocycles

Thomas Traynor
Illinois Wesleyan University

Rebecca Roesner, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Traynor, Thomas and Roesner, Faculty Advisor, Rebecca, "Host-Guest Interactions Between Niobium Polyoxometalates and Neutral Aza-Macrocycles" (2008). *John Wesley Powell Student Research Conference*. 25.

<https://digitalcommons.iwu.edu/jwprc/2008/posters2/25>

This is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P48

**HOST-GUEST INTERACTIONS BETWEEN NIOBIUM POLYOXOMETALATES
AND NEUTRAL AZA-MACROCYCLES**

Thomas Traynor and Rebecca Roesner*
Chemistry Department, Illinois Wesleyan University

During recent semesters, Dr. Roesner's research group has been investigating interactions between aza-macrocycles and polyoxometalates with the long-term goal of using polyoxometalates to direct the assembly of those macrocycles into supramolecular species. During those initial investigations, the group focused on Keggin polyoxoanions (e.g. $\text{PW}_{12}\text{O}_{40}^{3-}$, $\text{SiW}_{12}\text{O}_{40}^{4-}$), which are stable only at low pH conditions that also lead to protonation of the aza-macrocycles. The resulting polyoxoanion/ macrocycle adducts had properties consistent with the strong ionic bonds that held them together. They were poorly soluble in both water and polar organic solvents. In an effort to develop more-soluble polyoxoanion/adducts, the synthesis of niobium polyoxometalates (e.g. $\text{Nb}_6\text{O}_{19}^{7-}$ and $\text{Nb}_6\text{O}_{19}^{6-}$) is now underway. These niobium polyoxometalates are stable in basic solution where it will be possible to observe their interactions with neutral (free base) aza-macrocycles. It is hoped that the weaker, ion-dipole forces present in these systems will allow for greater solubility. This solubility should, in turn, allow for more-thorough spectroscopic investigation and the possibility of carrying out solution-based organic transformations on the polyoxoanion/macrocycle adducts.