



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2006, 17th Annual JWP Conference

Apr 8th, 9:00 AM - 10:00 AM

Quantifiable Torques for Microfluidic Studies

John Lattyak
Illinois Wesleyan University

Gabriel Spalding, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Lattyak, John and Spalding, Faculty Advisor, Gabriel, "Quantifiable Torques for Microfluidic Studies" (2006). *John Wesley Powell Student Research Conference*. 19.
<https://digitalcommons.iwu.edu/jwprc/2006/posters/19>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P41

QUANTIFIABLE TORQUES FOR MICROFLUIDIC STUDIES

John Lattyak and Gabriel Spalding*
Physics Department, Illinois Wesleyan University

Huge scientific advances have arisen from the ability to measure forces to high resolution. While Atomic Force Microscopes can reliably measure forces in solution down to around 10 pico-Newtons, force measurements well below that scale have been required to elucidate, e.g., basic mechanisms involved in the molecular motors responsible for muscle response. In this regime, non-invasive optical forces have become an invaluable tool. Just this year, the range of convincing calibration of optical forces has been extended down to 20 femto-Newtons! This opens up, for example, the possibility of studying the entropic forces involved in the statistical mechanics of DNA.

The study of torques has received less attention. It is our intention to use optical forces to push the limits of sensitivity for the measurement of torque in solution.

Our initial efforts have involved reproducing the current state of the art, an approach developed in 2004 by the Queensland group.¹ We utilize a circularly polarized laser beam to transfer angular momentum to a birefringent „probe“ particle (a micro-sphere of CaCO_3) in solution. By tightly focusing the laser beam, we create strong electromagnetic gradients that allow us to trap and manipulate the probe. Our ability to quantify the torques generated by our laser beam will depend on our knowledge of three key parameters: the frequency of the laser light, the power applied to the probe particle, and changes (per photon) in the polarization state of the transmitted beam.

1. „Optical Microrheology Using Rotating Laser-Trapped Particles,“ A. Bishop, T. Nieminen, N. Heckenberg, H. Runbinstein-Dunlop, Phys. Rev. Lett. 92, 198104 (2004).