



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2006, 17th Annual JWP Conference

Apr 8th, 9:00 AM - 10:00 AM

Using Fluorescence to Monitor Structural Changes in the Dimerization Helix of the Transcription Factor FNR

Kate McCulloch
Illinois Wesleyan University

Laura Moore, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

McCulloch, Kate and Moore, Faculty Advisor, Laura, "Using Fluorescence to Monitor Structural Changes in the Dimerization Helix of the Transcription Factor FNR" (2006). *John Wesley Powell Student Research Conference*. 23.
<https://digitalcommons.iwu.edu/jwprc/2006/posters/23>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P49

USING FLUORESCENCE TO MONITOR STRUCTURAL CHANGES IN THE DIMERIZATION HELIX OF THE TRANSCRIPTION FACTOR FNR

Kate McCulloch and Laura Moore*
Chemistry Department, Illinois Wesleyan University

Fumarate nitrate reductase (FNR) is an oxygen-sensing transcription factor in the facultative anaerobic bacteria *E. coli*. Although FNR is a widely-studied protein, little is known about its structure. FNR is a sequence homologue to catabolite receptor protein (CRP) in *E. coli*, and its structure is likely similar to CRP. When oxygen is absent, FNR is in an active dimeric form and contains a [4Fe-4S] cluster. Upon exposure to oxygen, the [4Fe-4S] cluster degrades to a [2Fe-2S] cluster and the protein subunits dissociate to two monomers. The structural changes that occur between these two forms are not well characterized. In this study, we use fluorescence spectroscopy to better understand the conformational changes between the active and inactive forms of FNR. Tryptophan, an amino acid with intrinsic fluorescent properties, has been used to replace some amino acid residues within the proposed dimerization helix of FNR. Changes in fluorescence emission wavelength indicate changes in the environment around the tryptophan, and are used to determine relative positions of amino acid residues within the dimerization helix. From preliminary data, the mutant FNR protein with tryptophan at position G149 does not have any significant change in emission wavelength upon exposure to oxygen, suggesting the environment around this residue does not change. However, the emission wavelength of the mutant protein at position M143 differs significantly between the anaerobic and aerobic forms. These data suggest that position 143 becomes exposed to solvent in the monomeric form and is isolated from solvent in the active dimeric form of the protein.