



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2006, 17th Annual JWP Conference

Apr 8th, 2:35 PM - 3:35 PM

Chromatographic Analysis of Pisum Sativum Porphobilinogen Synthase Morpheesins

Lindsay Sicks
Illinois Wesleyan University

David Bollivar, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Sicks, Lindsay and Bollivar, Faculty Advisor, David, "Chromatographic Analysis of Pisum Sativum Porphobilinogen Synthase Morpheesins" (2006). *John Wesley Powell Student Research Conference*. 1.

<https://digitalcommons.iwu.edu/jwprc/2006/oralpres3/1>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster and Oral Presentations P70, O3.1

**CHROMATOGRAPHIC ANALYSIS OF PISUM SATIVUM
PORPHOBILINOGEN SYNTHASE MORPHEEINS**

Lindsay Sicks and David Bollivar*

Biology Department, Illinois Wesleyan University

Porphobilinogen synthase (PBGS) is a metalloenzyme present in all organisms that synthesize tetrapyrrole compounds such as heme, chlorophyll, and vitamin B-12. PBGS catalyzes the condensation of two molecules of 5-aminolevulinic acid to form the tetrapyrrole precursor porphobilinogen. An artificial gene encoding PBGS of pea (*Pisum sativum*) has previously been designed, expressed, and purified (Kervinen, et al., *Biochemistry* 39:9018-9029). The specific activity is protein concentration dependent, which indicates that a maximally active enzyme can dissociate into less active smaller units. It has been hypothesized, and subsequently supported, that pea PBGS, essential for chlorophyll biosynthesis, exists as an equilibrium of an active octamer and an inactive hexamer. These different quaternary isoforms have been named "morpheins." Using a gel filtration (sizing) column, attempts to separate and characterize the morpheins of pea PBGS were made. Trials were carried out in various buffers, in which the pea protein ran at a characteristic octameric elution time. Further morphein separation was achieved on an anion exchange column, where separation of putative octamer and hexamer peaks was observed and the ratio of octamer to hexamer was directly related to the concentration of PBGS. Peaks were analyzed multiple ways, leading to the conclusions that the presence of Mg^{2+} ions appears to stabilize the octameric form of the pea PBGS, and the absence of Mg^{2+} ions appears to pull the dynamic equilibrium toward the hexameric form. These are the first analyses of the equilibrium mixture of pea quaternary forms under assay conditions. Further study is needed to confirm any interpretations.