



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2001, 12th Annual JWP Conference

Apr 21st, 11:00 AM - 12:30 PM

Systematic Relationships of Natalid and Furipterid Bats, Based on Hyoid Morphology (Chiroptera: Natalidae and Furipteridae)

Kosha Baxi
Illinois Wesleyan University

Thomas Griffiths, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Baxi, Kosha and Griffiths, Faculty Advisor, Thomas, "Systematic Relationships of Natalid and Furipterid Bats, Based on Hyoid Morphology (Chiroptera: Natalidae and Furipteridae)" (2001). *John Wesley Powell Student Research Conference*. 2.
<https://digitalcommons.iwu.edu/jwprc/2001/oralpres2/2>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation 2.3

**SYSTEMATIC RELATIONSHIPS OF NATALID AND FURIPTERID BATS,
BASED ON HYOID MORPHOLOGY
(CHIROPTERA: NATALIDAE AND FURIPTERIDAE)**

Kosha Baxi and Thomas Griffiths*

Department of Biology, Illinois Wesleyan University

The musculature of the hyoid region of two species of bats (Order Chiroptera) from two separate families, Natalidae and Furipteridae, was examined via microscopic dissection. Different morphological characters were scored and then entered into the computer program PAUP (Phylogenetic Analysis Using Parsimony), along with characters of families previously examined by Griffiths. A cladogram was generated. The analysis revealed support for the placement of Natalidae and Furipteridae together within the Superfamily Nataloidea, along with the families Thyropteridae and Myzopodidae. This grouping of Myzopodidae is quite surprising from a geographical standpoint. Myzopodids are endemic only to Madagascar, while thyropterids, natalids and furipterids are found in Central America and northern South America. Thus, the placement of Myzopodidae with the other three geographically close families implies that all four of these families share an unknown common ancestor in Africa.