



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2001, 12th Annual JWP Conference

Apr 21st, 11:00 AM - 12:30 PM

The Zoomin' HOTs: Dynamic Holographic Optical Tweezing

M. Bradley Taylor
Illinois Wesleyan University

Alexander J. Laurie
Illinois Wesleyan University

Gabriel C. Spalding, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Taylor, M. Bradley; Laurie, Alexander J.; and Spalding, Faculty Advisor, Gabriel C., "The Zoomin' HOTs: Dynamic Holographic Optical Tweezing" (2001). *John Wesley Powell Student Research Conference*. 4.

<https://digitalcommons.iwu.edu/jwprc/2001/oralpres2/4>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation 2.4

THE ZOOMIN' HOTs: DYNAMIC HOLOGRAPHIC OPTICAL TWEEZING

M. Bradley Taylor and Alexander J. Laurie and Gabriel C. Spalding*

Department of Physics, Illinois Wesleyan University

A new experimental technique, Holographic Optical Tweezing (H.O.T.), has been developed through a collaboration between our group at IWU and researchers at the University of Chicago.¹ Our addition of diffractive optics to earlier techniques for optical trapping significantly enhances the ability to control the assembly of biological cells or microscopic dielectric particles. In this new work, we describe original designs for optical systems using static holograms that nevertheless allow dynamic control over the number of traps in an assembly, their configuration/disorder (via spatial filtering), and even their spacings (via carefully designed zoom optics), as well as the strength of the traps (via the laser intensity). Although there are many possible applications for this new technology, as physicists our first interest is in exploring collective effects in optical binding, and in creating model systems that afford us many tunable parameters with which to explore many-body interactions. We are also interested in the onset of symmetry-breaking behavior (e.g., avalanches and the possibility of self-organized criticality, and the microscopic mechanisms underlying phase transitions in two and three dimensions).

¹Eric R. Dufresne, Gabriel C. Spalding, Matthew T. Dearing, Steven A. Sheets, David G. Grier, Review of Scientific Instruments 72, 1810 (2001).