



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2001, 12th Annual JWP Conference

---

Apr 21st, 10:00 AM - 11:00 AM

## Testing the Effects of Food Density on Echinoderm Larval Growth and the Correlation Between Juvenile Biomass and Size

Matthew Schnabl, '01  
*Illinois Wesleyan University*

William Jaeckle, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Schnabl, '01, Matthew and Jaeckle, Faculty Advisor, William, "Testing the Effects of Food Density on Echinoderm Larval Growth and the Correlation Between Juvenile Biomass and Size" (2001). *John Wesley Powell Student Research Conference*. 7.  
<https://digitalcommons.iwu.edu/jwprc/2001/posters/7>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation 19

**TESTING THE EFFECTS OF FOOD DENSITY ON ECHINODERM LARVAL  
GROWTH AND THE CORRELATION BETWEEN  
JUVENILE BIOMASS AND SIZE**

Matthew Schnabl and William Jaeckle\*

Department of Biology, Illinois Wesleyan University

I tested the hypothesis that juvenile size (surface area) accurately predicts juvenile organic content (biomass) in the sea urchin *Arbacia punctulata* (Echinodermata). Results of previous experiments with sea urchins suggest that juvenile size and survivorship are directly related to size of the eggs and free-swimming developmental stages (larvae) and the amount of available food (Emlet et al., 1987). To date, there is no published work that tests the hypothesis assumption that a larger juvenile possesses more biomass. If juvenile size does not predict juvenile biomass, then size per se may not be the critical factor predicting juvenile survivorship. Results of research presented last year revealed that when larvae were provided with an unlimited amount of food, juvenile size (surface area) significantly effects the amount of organic content, but there was not a predictive pattern between the two. I have extended this research project to evaluate the influence of a specific diet ration on juvenile size, biomass, and the correlation between these two variables. To date, after 10 weeks of continuous culturing, no larvae reared at the low food density ( $5 \times 10^2$  cells per ml) completed development to the juvenile stage. In contrast, 82 larvae in the high food treatment ( $5 \times 10^3$  cells per ml) successfully metamorphosed to become juvenile sea urchins. This experiment is ongoing and measurements of juvenile size, biomass, and their correlation are pending.