



Apr 14th, 9:00 AM - 10:00 AM

Fluid Flow Through Lytechinus Variegatus Sea Urchin Larvae

Sarah Lewis, '07

Illinois Wesleyan University

William Jaeckle, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Lewis, '07, Sarah and Jaeckle, Faculty Advisor, William, "Fluid Flow Through Lytechinus Variegatus Sea Urchin Larvae" (2007). *John Wesley Powell Student Research Conference*. 29.

<https://digitalcommons.iwu.edu/jwprc/2007/posters/29>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P53

FLUID FLOW THROUGH LYTECHINUS VARIEGATUS SEA URCHIN LARVAE

Sarah Lewis and William Jaeckle*

Biology Department, Illinois Wesleyan University

Sea urchins (Echinodermata: Echinoidea) produce developmental stages called larvae that feed and develop within the water column prior to becoming a bottom-dwelling juvenile. We evaluated the ability of larvae of the sea urchin *Lytechinus variegatus* to assimilate dissolved organic material (DOM) from seawater using fluorescence microscopy. In particle-free seawater, the larval digestive system readily absorbed macromolecules (a protein and a polysaccharide, 1mg/mL). With continued exposure, the label was detected within the body cavity suggesting that these molecules were distributed from the digestive system to other areas. To assess the effect of DOM on rates of particle capture, larvae were exposed to polystyrene beads (26,046/mL, 3 μ m diameter) in the presence or absence of DOM (the protein bovine serum albumin (BSA)). Rates of particle capture by larvae were affected by the presence of BSA, but the larvae of different ages responded differently. Younger larvae captured more beads in the absence of BSA while older larvae captured more beads in the presence of BSA.