



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2007, 18th Annual JWP Conference

Apr 14th, 9:00 AM - 10:00 AM

Social Ostracism and the Effect on Electroencephalogram Activity

Genevieve Nehrt, '07
Illinois Wesleyan University

Joseph Williams, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Nehrt, '07, Genevieve and Williams, Faculty Advisor, Joseph, "Social Ostracism and the Effect on Electroencephalogram Activity" (2007). *John Wesley Powell Student Research Conference*. 26.

<https://digitalcommons.iwu.edu/jwprc/2007/posters/26>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P59

SOCIAL OSTRACISM AND THE EFFECT ON ELECTROENCEPHALOGRAM ACTIVITY

Genevieve Nehrt and Joseph Williams*

Psychology Department, Illinois Wesleyan University

Previous fMRI research on social ostracism during participation in a ball-tossing task found increased activity in the anterior cingulate cortex when participants were not included in the ball-tossing game. This study aims to expand upon this previous research by seeing if similar results can be observed using EEG technology in a more realistic behavioral task, such as a chat room environment. This paradigm allows researchers to better understand the effects of social rejection on psychological, behavioral, and physiological responses and allows the participant more freedom of movement. It is hypothesized that when participants are excluded from the conversation, there will be an increase in theta EEG activity in the prefrontal cortex and less activation once the participants have been re-included into the conversation.