



Apr 14th, 9:00 AM - 10:00 AM

The Phylogenetic Distribution of Porphobilinogen Synthase (PBGS) Among Invertebrate Phyla: An Investigation of the Presence of PBGS in *Nematostella Vectensis*

Kelly Samartino '08
Illinois Wesleyan University

David Bollivar, Faculty Advisor
Illinois Wesleyan University

William Jaeckle, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Samartino '08, Kelly; Bollivar, Faculty Advisor, David; and Jaeckle, Faculty Advisor, William, "The Phylogenetic Distribution of Porphobilinogen Synthase (PBGS) Among Invertebrate Phyla: An Investigation of the Presence of PBGS in *Nematostella Vectensis*" (2007). *John Wesley Powell Student Research Conference*. 21.
<https://digitalcommons.iwu.edu/jwprc/2007/posters/21>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P71

**THE PHYLOGENETIC DISTRIBUTION OF PORPHOBILINOGEN SYNTHASE (PBGS)
AMONG INVERTEBRATE PHyla: AN INVESTIGATION OF THE PRESENCE OF PBGS IN
NEMATOSTELLA VECTENSIS**

Kelly Samartino and David Bollivar* and William Jaeckle*
Biology Department, Illinois Wesleyan University

The enzyme porphobilinogen synthase (PBGS) is known to catalyze the biosynthesis of tetrapyrroles such as heme and chlorophyll. The PBGS enzyme has been found in microorganisms, plants, vertebrate and invertebrate animals. This study was an investigation of the phylogenetic distribution of PBGS among invertebrate phyla. There is one example of an animal that lacks the PBGS enzyme, *Caenorhabditis elegans* (phylum Nematoda), thereby necessitating the acquisition of tetrapyrroles from its diet. Using a standard assay for PBGS, the presence of the enzyme was confirmed in the photosynthetic bacterium *Rhodobacter capsulatus*. No evidence of PBGS activity was demonstrated in *Nematostella vectensis*, a sea anemone (phylum Cnidaria); although, it was established that the *Nematostella vectensis* contained heme using a qualitative pyridine hemochromogen method.