



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2007, 18th Annual JWP Conference

---

Apr 14th, 2:35 PM - 3:35 PM

## **The Assimilation and Distribution of Dissolved Organic Matter By Planulae of Sea Anemones**

Amy Slott, '07  
*Illinois Wesleyan University*

William Jaeckle, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Slott, '07, Amy and Jaeckle, Faculty Advisor, William, "The Assimilation and Distribution of Dissolved Organic Matter By Planulae of Sea Anemones" (2007). *John Wesley Powell Student Research Conference*. 21.

<https://digitalcommons.iwu.edu/jwprc/2007/posters2/21>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P74

**THE ASSIMILATION AND DISTRIBUTION OF DISSOLVED ORGANIC MATTER  
BY PLANULAE OF SEA ANEMONES**

Amy Slott and William Jaeckle\*

Biology Department, Illinois Wesleyan University

The free-living developmental stage of sea anemones (phylum Cnidaria) is the planula larva. In general the feeding biology of planulae is poorly known and there is only one report of the ability of planulae to assimilate dissolved organic materials (DOM, specifically amino acids) from seawater (Ben-David-Zaslow and Benayahu, 2000). We examined the ability of planula larvae to assimilate dissolved proteins and polysaccharides from seawater and then distribute these materials throughout their bodies. Planulae were incubated, for known time intervals, in seawater containing ferritin, iron dextran, or bovine serum albumin (1mg / mL) and examined using light and fluorescence microscopy. With longer exposure times there was an increase in the amount of label in only the larval pharynx and endoderm; there was no evidence that these materials were distributed to the overlying ectoderm. Our results reveal a potential source of nutrition for the development of planula larvae.