



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2007, 18th Annual JWP Conference

Apr 14th, 2:35 PM - 3:35 PM

Effects of Dephosphorylation In 7E Mutants of NPR-A

Bridget Wall, '08

Illinois Wesleyan University

Brian Walter, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Wall, '08, Bridget and Walter, Faculty Advisor, Brian, "Effects of Dephosphorylation In 7E Mutants of NPR-A" (2007). *John Wesley Powell Student Research Conference*. 2.
<https://digitalcommons.iwu.edu/jwprc/2007/posters2/2>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P3, P82

EFFECTS OF DEPHOSPHORYLATION IN 7E MUTANTS OF NPR-A

Bridget Wall and Brian Walter*

Biology Department, Illinois Wesleyan University

Natriuretic peptides (designated ANP, BNP, and CNP) are hormones and paracrine factors which regulate blood volume, blood pressure, ventricular hypertrophy, pulmonary hypertension, fat metabolism and long bone growth. When bound to natriuretic peptide receptors, they allow for the production of cGMP (cyclic guanosine monophosphate), a secondary messenger which allows for the production of proteins from the cell. These receptors are designated NPR-A/GC-A and NPR-B/GC-B. Both are guanylyl cyclases (enzymes that synthesize cGMP). The focus of this research is to further examine the involvement of dephosphorylation in the desensitization of natriuretic peptide receptors. Previously, phosphorylation sites on the guanylyl cyclase domain have been mutated to express glutamate, an amino acid with a charge of -1 (the charge of a phosphate is -2). Therefore, the glutamate is mimicking the charge on the phosphorylated residue present when the receptor is phosphorylated.

Ergo, these NPR-A mutants (hereafter referred to as 7E mutants) are constitutively phosphorylated and always active. Within the current model, there should be a significant difference in cGMP production when stimulated with natriuretic peptides. By continuing previous studies geared toward a better understanding of phosphorylation in these receptors, members of the field of pharmacology will be able to develop a drug that will either semi-permanently dephosphorylate or phosphorylate natriuretic peptide receptors in an effort to treat heart disease and hypertension.