



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2002, 13th Annual JWP Conference

Apr 21st, 1:15 PM - 2:15 PM

Abiotic Factors Influencing Body Color in *Murgantia Histrionica*

Jennifer Olson, '02
Illinois Wesleyan University

Given Harper, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Olson, '02, Jennifer and Harper, Faculty Advisor, Given, "Abiotic Factors Influencing Body Color in *Murgantia Histrionica*" (2002). *John Wesley Powell Student Research Conference*. 7.

<https://digitalcommons.iwu.edu/jwprc/2002/posters3/7>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P40

**ABIOTIC FACTORS INFLUENCING BODY COLOR IN
*MURGANTIA HISTRIONICA***

Jennifer Olson and Given Harper*

Department of Biology, Illinois Wesleyan University

Many insects have developed the ability to thermoregulate by basking in the sun in order to maintain a higher body temperature. Because dark colors absorb more solar radiation, some insects will also alter their body color in colder weather in order to thermoregulate more efficiently. This experiment was designed to determine the effects of thermoperiod and photoperiod on the body color of the harlequin bug, *Murgantia histrionica*. Two populations were raised from the nymph stage. One treatment was exposed to simulated summer conditions with a 15:9hr, light:dark (L:D) photoperiod, and 30:24°C (day:night) thermoperiod, while the other treatment was exposed to simulated fall conditions with a 9:15hr L:D photoperiod and 22:15°C thermoperiod. After one generation, adults were examined for differences in the ratios of black to yellow body color. As predicted, the harlequin bugs in the first treatment had significantly more yellow color on their dorsal side than those in the second treatment ($t=4.67$, $df=9$, $p<.000$). The increase of black color in the fall population is most likely facilitating their absorption of more solar radiation. This allows them to compensate for the lower temperature and the shorter duration of daily light.