



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2009, 20th Annual JWP Conference

---

Apr 18th, 10:00 AM - 11:00 AM

## Biochemical Characterization of the *Candida Albicans* PBGS Enzyme

Bryan Rea, '09  
*Illinois Wesleyan University*

David Bollivar, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Rea, '09, Bryan and Bollivar, Faculty Advisor, David, "Biochemical Characterization of the *Candida Albicans* PBGS Enzyme" (2009). *John Wesley Powell Student Research Conference*. 2.

<https://digitalcommons.iwu.edu/jwprc/2009/oralpres2/2>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Oral Presentation O2.2

**BIOCHEMICAL CHARACTERIZATION OF THE  
*CANDIDA ALBICANS* PBGS ENZYME**

Bryan Rea and David Bollivar\*

Biology Department, Illinois Wesleyan University

The enzyme porphobilinogen synthase (PBGS) catalyzes the reaction that forms porphobilinogen, an important precursor in the biosynthetic pathways that lead to the production of heme, chlorophyll, and other tetrapyrroles. Previous studies have indicated that PBGS often exists in two states, termed morphoforms, an active octameric form and inactive hexameric form. Recent evidence has shown that certain molecules may be able to lock the enzyme as a hexamer, thereby inactivating it. Such selective agents against the PBGS enzyme could be very effective in dealing with pathogenic organisms. One such pathogen is *Candida albicans*, a fungus responsible for thrush, most common in babies and immunocompromised individuals, e.g. those with AIDS. The goal of our current study was to isolate, purify, and characterize the PBGS enzyme from *C. albicans*. Our data indicate that our purified product is indeed capable of catalyzing the reaction to form porphobilinogen. Furthermore, our results indicate that the optimum pH of the enzyme is approximately 8.7, and it requires the presence of zinc and reducing agents. Further study will shed light on the optimal level of zinc required, other metal ion requirements (such as magnesium), and identification of possible inhibitors of enzymatic activity. The results obtained will provide a much clearer picture of the general biochemistry of the *C. albicans* PBGS. Such information combined with structural studies, like NMR spectroscopy or X-ray crystallography, may lead to the development of an agent that can selectively inhibit the *C. albicans* PBGS enzyme.