



Apr 18th, 2:35 PM - 3:35 PM

## The Expression and Function of TRPV1 and TRPV3 in the Hippocampus

Kathleen Louis  
*Illinois Wesleyan University*

Given Harper, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Louis, Kathleen and Harper, Faculty Advisor, Given, "The Expression and Function of TRPV1 and TRPV3 in the Hippocampus" (2009). *John Wesley Powell Student Research Conference*. 12.

<https://digitalcommons.iwu.edu/jwprc/2009/posters2/12>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P26

**THE EXPRESSION AND FUNCTION OF TRPV1 AND TRPV3  
IN THE HIPPOCAMPUS**

Kathleen Louis and Given Harper\*

Biology Department, Illinois Wesleyan University and  
Department of Pharmacology, Southern Illinois School of Medicine

Sensory neurons in the peripheral nervous system are responsible for sending sensory information from the periphery to the central nervous system. The Transient Receptor Potential Vanilloid (TRPV) family of genes encodes for different types of channels in the peripheral nervous system that respond to temperature, mechanical and chemical stimuli. Particularly, TRPV1 is expressed in sensory neurons and TRPV3 is expressed in keratinocytes (skin cells). But it is becoming apparent that the role of TRPV channels is much broader than just sensory transduction because they are proposed to be expressed in certain regions of the brain. The purpose of this study was to determine the expression and function of TRPV1 and TRPV3 in the hippocampus of the brain. In this study, using patch clamp electrophysiological techniques, I recorded membrane currents and membrane depolarizations leading to the generation of action potentials from cultured hippocampal neurons in response to application of 2-APB, a TRPV3 channel activator. I also used the calcium fluorescence imaging technique and found that application of 2-APB induced an increase in intracellular  $\text{Ca}^{2+}$  levels. However, application of capsaicin, a TRPV1 activator, neither induced a current nor increased intracellular  $\text{Ca}^{2+}$  levels in the same experimental conditions. From these results, I conclude that TRPV3 channels are present in the hippocampus, but not TRPV1 channels.