



Apr 18th, 2:35 PM - 3:35 PM

## Assembly of Niobium Polyoxometalates and Azamacrocycles

Christopher J. Santee  
*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Santee, Christopher J. and Roesner, Faculty Advisor, Rebecca, "Assembly of Niobium Polyoxometalates and Azamacrocycles" (2009). *John Wesley Powell Student Research Conference*. 23.

<https://digitalcommons.iwu.edu/jwprc/2009/posters2/23>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P48

**ASSEMBLY OF NIOBIUM POLYOXOMETALATES AND AZAMACROCYCLES**

Christopher J. Santee and Rebecca Roesner\*  
Chemistry Department, Illinois Wesleyan University

Dr. Roesner's research group has been investigating interactions between azamacrocycles and polyoxometalates with the long-term goal of using polyoxometallates to direct the assembly of those macrocycles into supramolecular species. Initially, the group focused on Keggin polyoxoanions (e.g.  $\text{PW}_{12}\text{O}_{40}^{3-}$ ,  $\text{SiW}_{12}\text{O}_{40}^{4-}$ ), which are stable only in acidic solution. The azamacrocycles are partially or fully protonated over this pH range and the resulting polyoxoanion/macrocycle adducts exhibit properties consistent with their strong ionic bonding. Most of the salts composed of macrocyclic cations and polyoxoanions are poorly soluble in both water and polar organic solvents. In an effort to develop more-soluble polyoxoanion/adducts, the hexaniobate ion,  $\text{Nb}_6\text{O}_{19}^{8-}$ , was synthesized. This niobium polyoxometallate is stable at high pH and can therefore coexist in solution with neutral (free base) azamacrocycles. To date, no host guest complexes between the hexaniobate ion and free base azamacrocycles have been isolated.