



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2009, 20th Annual JWP Conference

---

Apr 18th, 9:00 AM - 10:00 AM

**The Effect of Predominant Light on Coloration in the Chinese  
Praying Mantid *Tenodera aridifolia sinensis***

Lukasz Sewera  
*Illinois Wesleyan University*

Elizabeth Balser, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Sewera, Lukasz and Balser, Faculty Advisor, Elizabeth, "The Effect of Predominant Light on Coloration in the Chinese Praying Mantid *Tenodera aridifolia sinensis*" (2009). *John Wesley Powell Student Research Conference*. 25.

<https://digitalcommons.iwu.edu/jwprc/2009/posters/25>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P49

**THE EFFECT OF PREDOMINANT LIGHT ON COLORATION IN THE CHINESE  
PRAYING MANTID *TENODERA ARIDIFOLIA SINENSIS***

Lukasz Sewera and Elizabeth Balser\*  
Bioogy Department, Illinois Wesleyan University

*Tenodera aridifolia sinensis*, also known as the Chinese praying mantid, is an introduced species in North America. With no major predators, this insect has thrived and successfully incorporated itself into ecosystems throughout the Midwestern and Eastern United States. Juveniles of *T. aridifolia sinensis* are easy to rear in laboratory settings, making it an ideal insect for experimentation. Specimens of *T. aridifolia sinensis* have three predominant morphological patterns with respect to exoskeleton coloration; they are green, brown, or a mixture of the two. As such, these insects are usually found in habitats similar to their color (i.e., green mantids are found on leaves, brown ones on bark) (Prete 1999). In this study, we sought to determine whether mantid exoskeleton color is influenced by the predominating color of light in which they live. Approximately 100 nymphs of *T. aridifolia sinensis* were reared from birth, isolated in chambers with colored light (red, blue, green, and clear). All other environmental factors were standardized. Color was quantitatively assessed using a spectrophotometer to determine the concentration of the pigment Biliverdin IXa in each specimen. Biliverdin IXa is responsible for the green coloration in praying mantids (Rudiger 1968). Differences in color could be due to varying concentrations of this pigment resulting from the effect of predominant light.