



Apr 10th, 10:00 AM - 11:00 AM

The Uptake of Dissolved Molecules by the Freshwater Rotifer, *Asplanchna Priodonta*

Stephanie R. Ross
Illinois Wesleyan University

William Jaeckle, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Biology Commons](#)

Ross, Stephanie R. and Jaeckle, Faculty Advisor, William, "The Uptake of Dissolved Molecules by the Freshwater Rotifer, *Asplanchna Priodonta*" (2010). *John Wesley Powell Student Research Conference*. 3.

<https://digitalcommons.iwu.edu/jwprc/2010/oralpres/3>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation O1.2

**THE UPTAKE OF DISSOLVED MOLECULES BY THE
FRESHWATER ROTIFER, *ASPLANCHNA PRIODONTA***

Stephanie R. Ross and William Jaeckle*
Biology Department, Illinois Wesleyan University

DOM (dissolved organic matter) is an abundant potential source of nutrition for aquatic invertebrates, however, its nutritional significance is not well documented. Rotifers are water aquatic metazoans which utilize a ciliated corona for food acquisition. The carnivorous freshwater rotifer, *Asplanchna priodonta*, was exposed to labeled proteins and polysaccharides at substrate concentrations of $\leq 1\text{mg/mL}$ for 1-18 hours. Initially, the labels were observed within the stomach lumen and gastric epithelium. With extended exposure, the intensity of the label increased and was detected within the body cavity. These results suggest that the gastric cavity is continually exposed to freshwater and DOM present may be absorbed. This is consistent with previous studies describing digestion as initially extracellular and subsequently intracellular within endocytotic vesicles. The presence of labeled compounds within stomach cells and the body cavity of *A. priodonta* demonstrates this species's ability to digest and assimilate DOM as a possible source of nutrition.