



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2010, 21st Annual JWP Conference

Apr 10th, 9:00 AM - 10:00 AM

Variable Absorption of Dissolved Organic Material by Pilidium Larvae

Kristina Bennett
Illinois Wesleyan University

William Jaeckle, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Biology Commons](#)

Bennett, Kristina and Jaeckle, Faculty Advisor, William, "Variable Absorption of Dissolved Organic Material by Pilidium Larvae" (2010). *John Wesley Powell Student Research Conference*. 7.

<https://digitalcommons.iwu.edu/jwprc/2010/posters/7>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P5

**VARIABLE ABSORPTION OF DISSOLVED ORGANIC MATERIAL
BY PILIDIUM LARVAE**

Kristina Bennett and William Jaeckle*
Biology Department, Illinois Wesleyan University

Many marine nemerteans develop through a free-swimming stage called a pilidium larva. Pilidia are known to feed on suspended single-celled algae and bacteria. Pilidia may use dissolved organic materials (DOM) in seawater as an additional source of nutrition. We investigated DOM absorption by pilidium larvae collected from two geographical regions. Larvae collected from the western Atlantic Ocean showed no detectable absorption of the protein ferritin (1 mg / mL) from seawater (≤ 10 h, 25°C). In contrast, when larvae of Antarctic nemertean *Parborlasia corrugatus* were exposed to the polysaccharide iron dextran (1 mg / mL) for ≤ 26 h (0°C) the label was detected in the cells of the digestive system. This pattern of absorption is explained by seawater flow into the digestive system. For tropical larvae, data do not support DOM as a potential nutrient source; while for polar larvae, it seems that DOM could serve as an augmenting resource.