



Apr 10th, 9:00 AM - 10:00 AM

## An Analysis of Continuous Chest Compression CPR for EMS Providers for Out of Hospital Cardiac Arrest

Megan Gleason  
*Illinois Wesleyan University*

Kira Hudson Banks, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Psychiatry and Psychology Commons](#)

---

Gleason, Megan and Hudson Banks, Faculty Advisor, Kira, "An Analysis of Continuous Chest Compression CPR for EMS Providers for Out of Hospital Cardiac Arrest" (2010). *John Wesley Powell Student Research Conference*. 13.

<https://digitalcommons.iwu.edu/jwprc/2010/posters/13>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P19

**AN ANALYSIS OF CONTINUOUS CHEST COMPRESSION CPR FOR EMS PROVIDERS FOR OUT OF HOSPITAL CARDIAC ARREST**

Megan Gleason and Kira Hudson Banks\*  
Psychology Department, Illinois Wesleyan University

Much research has been done to improve the outcomes of patients found in cardiac arrest outside the hospital. The American Heart Association has long advocated Advanced Cardiac Life Support (ACLS), a procedure that encompasses cycles of chest compressions with advanced airway maintenance and defibrillation. Recent evidence has suggested that these current guidelines are ineffective due to prolonged “hands off” time. New research suggests utilizing a technique known as continuous chest compression CPR that delays advanced airway management and instead focuses on defibrillation and continuous chest compressions. Across the country, research has demonstrated that when EMS providers utilize this technique and have support from the receiving hospital, survival to hospital discharge rates have increased from 4.7% (using standard ACLS protocols) to 17.6% (with the new technique). The Newark (OH) Fire Department protocols were modified to implement continuous chest compression CPR for the care of patients in cardiac arrest. The present research analyzes quality improvement (QI) / quality assurance (QA) data from this fire department to determine how the change in protocol affected patient outcome.