



Apr 10th, 2:35 PM - 3:35 PM

## Synthesis of Polyazamacrocyclic Molecular Tweezers as a Host For Polyoxoanions

Marie Sutor  
*Illinois Wesleyan University*

Ronald Grider  
*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

 Part of the [Chemistry Commons](#)

---

Sutor, Marie; Grider, Ronald; and Roesner, Faculty Advisor, Rebecca, "Synthesis of Polyazamacrocyclic Molecular Tweezers as a Host For Polyoxoanions" (2010). *John Wesley Powell Student Research Conference*. 27.

<https://digitalcommons.iwu.edu/jwprc/2010/posters2/27>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P52

## SYNTHESIS OF POLYAZAMACROCYCLE MOLECULAR TWEEZERS AS A HOST FOR POLYOXOANIONS

Marie Sutor, Ronald Grider and Rebecca Roesner\*  
Chemistry Department, Illinois Wesleyan University

Host-guest chemistry concerns the design of receptor molecules that complement their targeted guests with respect to size, shape, and charge. One common area in host-guest chemistry involves the design of molecular tweezers, which are shown schematically in Figure 1.




Figure 1. Schematic representation of molecular tweezers grasping a guest molecule.

Molecular tweezers have been developed to grasp a wide variety of neutral, cationic, and anionic guests. This study is directed toward the development of molecular tweezers for polyoxoanion guests. Because polyoxoanions are large, approximately spherical in shape, and negatively charged, they are complementary to the polyazamacrocyclic cations that will serve as pincers in the proposed molecular tweezers (Figure 2).




Figure 2. Proposed design of a molecular tweezers for polyoxoanion guests.

Toward these synthetic goals, the methods of Zhong, Mikkola, and Lönnberg are being employed in the multi-step synthesis of 1-ethylamino-1, 4, 7, 10, 13, 16, 19-heptaazacycloheicosane. Synthetic intermediates have been characterized by melting point and  $^1\text{H}$  NMR.