



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2011, 22nd Annual JWP Conference

Apr 9th, 9:00 AM - 10:00 AM

**Incidence of Larval Cloning in the Sea Urchins *Arbacia punctulata*
and *Lytechinus variegatus***

Lauren Knaack
Illinois Wesleyan University

William Jaeckle, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Biology Commons](#)

Knaack, Lauren and Jaeckle, Faculty Advisor, William, "Incidence of Larval Cloning in the Sea Urchins *Arbacia punctulata* and *Lytechinus variegatus*" (2011). *John Wesley Powell Student Research Conference*. 14.

<https://digitalcommons.iwu.edu/jwprc/2011/posters/14>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P27

INCIDENCE OF LARVAL CLONING IN THE SEA URCHINS *ARBACIA PUNCTULATA* AND *LYTECHINUS VARIEGATUS*

Lauren Knaack and William Jaeckle*
Biology Department, Illinois Wesleyan University

Cloning is the development of a new, genetically identical individual from a pre-existing organism. Cloning by free-living developmental forms called larvae has been observed in 4 of the 6 taxonomic classes within phylum Echinodermata. An earlier report by Vaughn and Strathmann (2008) indicated the rate of cloning by pluteus larvae of *Dendraster excentricus* is influenced by the surrounding environment. We tested this hypothesis by monitoring the incidence and frequency of cloning of larvae of the sea urchins *Arbacia punctulata* and *Lytechinus variegatus* in response to different sources and abundances of particulate foods. Groups of seven plutei were reared transferred into each of nine different shot glasses containing 40 mL 36 ‰ filtered seawater. Commercial food particles (EZ-larva®), or algal cells, were added to produce 3 food concentrations (low, medium, and high). Every two days the plutei were counted, fed, and transferred to a glass with filtered seawater and food. They were counted to see if any clones had been released and if any plutei were lost or died. Only two clonally produced larvae were produced by larvae of *L. variegatus*; *A. punctulata* larvae did not clone. Overall survivorship was directly related to abundance of food and ranged from 18% to 80%. The observed low rates of cloning ($\approx 2\%$ of cultured larvae) imply that cloning (1) is not a common developmental feature among larvae of sea urchins or (2) our environmental conditions were inappropriate to induce cloning. Cloning rates may differ in nature depending upon numerous factors not studied in these experiments such as water temperature, salinity, or concentration of larvae in culture.