



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2011, 22nd Annual JWP Conference

---

Apr 9th, 9:00 AM - 10:00 AM

## **Efficacy of a Cocaine-Schizophrenia Model in Rats: Behavioral and Physiological Measures**

Andrew W. Ross  
*Illinois Wesleyan University*

Joseph Williams, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Psychology Commons](#)

---

Ross, Andrew W. and Williams, Faculty Advisor, Joseph, "Efficacy of a Cocaine-Schizophrenia Model in Rats: Behavioral and Physiological Measures" (2011). *John Wesley Powell Student Research Conference*. 20.

<https://digitalcommons.iwu.edu/jwprc/2011/posters/20>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P41

**EFFICACY OF A COCAINE- SCHIZOPHRENIA MODEL IN RATS:  
BEHAVIORAL AND PHYSIOLOGICAL MEASURES**

Andrew W. Ross and Joseph Williams\*

Psychology Department, Illinois Wesleyan University

Behaviors exhibited during exposure to acute levels of cocaine are similar to behaviors exhibited by patients with schizophrenia and psychosis, leading researchers to hypothesize that rats exposed to acute levels of cocaine may serve as a valid research model for schizophrenia. In addition to behavior, cocaine addiction and schizophrenia share many other commonalities, including similar neurochemical and neuroanatomical substrates. This study looks to test the validity of this model by measuring whether cocaine exposure in rats results in common behavioral deficits (e.g., aggression, spatial memory and cognitive flexibility) observed in patients with schizophrenia. The behaviors were measured in a cocaine experimental group (5 mg/kg) and a saline control group. The aggression was measured based on the Resident-Intruder paradigm. Cognitive flexibility and spatial memory were measured by examining performance in an Attentional Set Shift paradigm. Though no statistical significance was found, the trends seen in the data for spatial memory and cognitive flexibility support the hypothesis. However, the results of the aggression measure do not support the hypothesis. Overall, more research is necessary to determine the overall validity of the use of cocaine exposed rats as a model of schizophrenia.