



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2011, 22nd Annual JWP Conference

---

Apr 9th, 2:35 PM - 3:35 PM

## Scanning Electrochemical Microscopic Imaging of Taste Cells

Aaron Moore  
*Illinois Wesleyan University*

Melinda Baur, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

 Part of the [Chemistry Commons](#)

---

Moore, Aaron and Baur, Faculty Advisor, Melinda, "Scanning Electrochemical Microscopic Imaging of Taste Cells" (2011). *John Wesley Powell Student Research Conference*. 17.  
<https://digitalcommons.iwu.edu/jwprc/2011/posters2/17>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P34

## **SCANNING ELECTROCHEMICAL MICROSCOPIC IMAGING OF TASTE CELLS**

Aaron Moore and Melinda Baur\*

Chemistry Department, Illinois Wesleyan University

The goal of this project is to detect the stimulated release of serotonin and to image the topography of taste cells using Scanning Electrochemical Microscopy (SECM). The sense of taste is important for animals because it allows animals to recognize food, derive pleasure from food, and to detect food that may be unsafe to eat. The multiple taste signaling pathways respond to five different kinds of taste molecules: sour, sweet, umami, salty and bitter. Taste buds are organized groups of 50-100 cells responsible for the transduction of the taste signal. Within the taste bud, there are at least three different cell types: Type I (Glial-like) Cells, Type II (Receptor) Cells and Type III (Pre-synaptic) Cells. While there is evidence that some taste transduction also occurs in Type I and Type III cells, currently the best understood signal transduction pathways occur in Type II cells. Type III cells contain serotonin and can depolarize in response to various stimuli. We can detect the release of electro-active neurotransmitters like serotonin, using electrochemistry. This type of electrochemical detection of released neurotransmitters has been used successfully with model neurons. The role of the Type III Cells in the in the taste signal transduction pathway can be elucidated using this new electrochemical tool. Taste bud extraction and isolation have been successfully completed at Illinois Wesleyan University and now the focus is to detect neurotransmitter release from isolated taste buds.