



Apr 9th, 2:35 PM - 3:35 PM

## Management of Permanent Pacing Wires During Tricuspid Valve Repair

Luke Sheen

*Illinois Wesleyan University*

Daniel L. Cherkassky, Faculty Advisor

Virna L. Sales, MD, Faculty Advisor

Richard Lee, MD, MBA, Faculty Advisor

Brittany Lapin, MPH, Faculty Advisor

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Medicine and Health Sciences Commons](#)

---

Sheen, Luke; Cherkassky, Faculty Advisor, Daniel L.; Sales, MD, Faculty Advisor, Virna L.; Lee, MD, MBA, Faculty Advisor, Richard; Lapin, MPH, Faculty Advisor, Brittany; Wang, PhD, Faculty Advisor, Edward; and McCarthy, MD, Faculty Advisor, Patrick M., "Management of Permanent Pacing Wires During Tricuspid Valve Repair" (2011). *John Wesley Powell Student Research Conference*. 26.

<https://digitalcommons.iwu.edu/jwprc/2011/posters2/26>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

---

### **Presenter and Advisor Information**

Luke Sheen; Daniel L. Cherkassky, Faculty Advisor; Virna L. Sales, MD, Faculty Advisor; Richard Lee, MD, MBA, Faculty Advisor; Brittany Lapin, MPH, Faculty Advisor; Edward Wang, PhD, Faculty Advisor; and Patrick M. McCarthy, MD, Faculty Advisor

Poster Presentation P44

## MANAGEMENT OF PERMANENT PACING WIRES DURING TRICUSPID VALVE REPAIR

Luke Sheen and Daniel L Cherkassky\*, Virna L. Sales\*, MD, Richard Lee\*, MD, MBA, †Brittany Lapin, MPH, †Edward Wang, PhD and Patrick M. McCarthy\*, MD, FACC  
Bluhm Cardiovascular Institute, \*Division of Cardiac Surgery, and †Clinical Trials Unit at Northwestern University Feinberg School of Medicine and Northwestern Memorial Hospital, Chicago, IL

**Objectives:** We compared 3 management techniques for permanent pacing wires by studying the recurrence of postoperative tricuspid regurgitation (TR).

**Methods:** From 2004-2009, 40 patients (mean age  $71 \pm 11$  years, 60% male) with prior permanent pacing wires in the right ventricle underwent tricuspid valve annuloplasty (TVA). Wires were managed with three different surgical approaches: leaving the wire in its transtricuspid position ( $n=22$ ); removal and epicardial replacement ( $n=10$ ); and securing it in the postero-septal commissure ( $n=8$ ). Compared to removed group, transtricuspid and secured groups had greater prevalence of pre-operative pulmonary hypertension ( $96\%[21/22]$  vs.  $88\%[7/8]$  vs.  $40\%[4/10]$ ,  $p=0.001$ ) and atrial fibrillation ( $86\%[19/22]$  vs.  $88\%[7/8]$  vs.  $40\%[4/10]$ ,  $p<0.05$ ). Ejection fraction was similar (transtricuspid 45%, removed 53%, and secured 47%;  $p=0.8$ ). The percentage of patients with NYHA class III/IV were similar among the 3 groups (transtricuspid  $77\%[17/22]$ , removed  $70\%[7/10]$ , and secured  $50\%[4/8]$ ;  $p=0.5$ ). Ambler score was highest in secured group ( $28 \pm 14$ ,  $p<0.05$ ) followed by transtricuspid ( $20 \pm 10$ ) and removed ( $14 \pm 12$ ) groups. A total of 148 preoperative, discharge, and late follow-up ( $>6$  months) transthoracic echocardiograms were analyzed. Mean follow-up was  $21.9 \pm 16.6$  months (range 10-1840 days) and complete in 28/29 (97%).

**Results:** One patient died in the hospital in each group after TVA. Incidence of TR 3+/4+ decreased in all groups at discharge ( $70\%[28/40]$  vs  $5\%[2/40]$ ,  $p=0.001$ ) and late follow-up ( $15\%[8/53]$ ,  $p<0.001$  with no 4+ TR) compared to preoperatively. Preoperatively, secured group had the highest significant TR compared with transtricuspid and removed groups ( $88\%[7/8]$  vs.  $68\%[15/22]$  vs.  $60\%[6/10]$ ,  $p=0.4$ ) but at discharge, only transtricuspid had TR recurrence ( $9\%[2/22]$ ,  $p=0.4$ ). At late follow-up, significant TR redeveloped and was lowest in secured and similar in removed and transtricuspid groups ( $14.3\%[1/7]$  vs.  $15.4\%[2/13]$  vs.  $15.2\%[5/33]$ ,  $p=0.998$ ).

**Conclusion:** Patients in secured group represent a cohort with high risk factors for TR recurrence (greater preoperative pulmonary hypertension, AF and high ambler score) compared with those in removed group, but similarly demonstrated decreased TR across time. High-risk transtricuspid patients showed worsening TR after repair. Securing pace wires into the commissure during TVR may be a reasonable approach in management of pacemaker wires in high risk patients.