



Apr 14th, 10:00 AM - 11:00 AM

**Comparison of Regional Eggshell Porosity Between the Brood
Parasitic Brown-Headed Cowbird (*Moluthrus ater*) and its Hosts:
the Dickcissel (*Spiza americana*), and Two Non-Parasitic
Relatives, the Red-Winged Blackbird (*Agelaius phoeniceus*) and
the Common Grackle (*Quiscalus quiscula*).**

Brittany Childs
Illinois Wesleyan University

W. B. Jaeckle, Faculty Advisor
Illinois Wesleyan University

R. G. Harper, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the **Biology Commons**
Illinois Wesleyan University

Childs, Brittany; Jaeckle, Faculty Advisor, W. B.; Harper, Faculty Advisor, R. G.; Kiefer, M.;
Owens, J. W., and Piller, B. D., "Comparison of Regional Eggshell Porosity Between the Brood
Parasitic Brown-Headed Cowbird (*Moluthrus ater*) and its Hosts: the Dickcissel (*Spiza
americana*), and Two Non-Parasitic Relatives, the Red-Winged Blackbird (*Agelaius
phoeniceus*) and the Common Grackle (*Quiscalus quiscula*).\" (2012). *John Wesley Powell
Student Research Conference*. 3.
<https://digitalcommons.iwu.edu/jwprc/2012/oralpres/3>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital
Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any
way that is permitted by the copyright and related rights legislation that applies to your use. For
other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights
are indicated by a Creative Commons license in the record and/ or on the work itself. This material
has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information,
please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Presenter and Advisor Information

Brittany Childs; W. B. Jaeckle, Faculty Advisor; R. G. Harper, Faculty Advisor; M. Kiefer; J. W. Rivers; and B. D. Peer

Oral Presentation O1.3

COMPARISON OF REGIONAL EGGSHELL POROSITY BETWEEN THE BROOD PARASITIC BROWN-HEADED COWBIRD (*MOLUTHRUS ATER*) AND ITS HOSTS: THE DICKCISSEL (*SPIZA AMERICANA*), AND TWO NON-PARASITIC RELATIVES, THE RED-WINGED BLACKBIRD (*AGELAIUS PHOENICEUS*) AND THE COMMON GRACKLE (*QUISCALUS QUISCULA*).

Brittany Childs^a, W.B. Jaeckle^{a*}, R.G. Harper^{a*}, M. Kiefer^a, J.W. Rivers^b, B.D. Peer^c

^aDepartment of Biology, Illinois Wesleyan University, Bloomington, IL

^bDepartment of Forest Ecosystems and Society, Oregon State University, Corvallis, OR

^cDepartment of Biological Sciences, Western Illinois University, Macomb, IL

The Brown-headed Cowbird is a generalist brood parasite that lays eggs in the nests of many host species, including the Dickcissel and two non-parasitic relatives: the Red-winged Blackbird and the Common Grackle. Cowbird eggs reportedly hatch sooner than equivalently-sized host eggs, presumably via accelerated embryonic development enabled by a greater eggshell porosity and consequently greater gas exchange. The distribution of pores among apical, equatorial and basal eggshell regions within cowbirds and host species is undetermined. We tested the hypothesis that equatorial porosity would be greatest because respiratory gases cross the eggshell and enter or exit the embryo's circulatory system in that region. We found that, when normalized to mass, cowbird eggshells had significantly greater pore area and porosity in equatorial regions compared to its three hosts ($p \leq 0.001$). Cowbird eggshells had greater apical pore area than the hosts ($p < 0.04$) but equivalent basal pore area ($p > 0.09$) and porosity ($p > 0.15$) to its relatives.