



Apr 14th, 9:00 AM - 10:00 AM

Isolation of *Rhodobacter capsulatus* Bacteriophages and development of Optimal Infection Conditions

Brooke Bernardoni
Illinois Wesleyan University

Matthew R. Bockman
Illinois Wesleyan University

David Bollivar, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

Bernardoni, Brooke; Bockman, Matthew R.; and Bollivar, Faculty Advisor, David, "Isolation of *Rhodobacter capsulatus* Bacteriophages and development of Optimal Infection Conditions" (2012). *John Wesley Powell Student Research Conference*. 6.
<https://digitalcommons.iwu.edu/jwprc/2012/posters/6>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P11

**ISOLATION OF *RHODOBACTER CAPSULATUS* BACTERIOPHAGES AND
DEVELOPMENT OF OPTIMAL INFECTION CONDITIONS**

Brooke Bernardoni, Matthew R. Bockman, and David W. Bollivar*
Biology Department, Illinois Wesleyan University

Few studies have been performed regarding bacteriophages that infect photosynthetic bacteria. *Rhodobacter capsulatus* is a photosynthetic bacterium that has been used as a model system for studying the genetics of photosynthesis. It has been used as a host for bacteriophages in the past, but most of this work was performed prior to the advent of molecular biology. Three bacteriophages have been isolated that will infect *Rhodobacter capsulatus*. During the course of the studies described in this poster, it was discovered that different strains of this bacterium have very different susceptibilities to infection by bacteriophages. The isolated bacteriophages were discovered from water samples taken from a creek in the Bloomington/Normal, Illinois area. The *R. capsulatus* strain YW1C-6 showed evidence for increased susceptibility for bacteriophage infection when compared to the St. Louis strain. Following isolation and purification of these bacteriophages, optimal conditions for bacteriophage infection were developed to enhance isolation techniques for further studies.