



Apr 14th, 2:35 PM - 3:35 PM

## Metabolic Contribution of Particulate Matter (<math>< \mu\text{m}</math>) for Larvae of *Lytechinus variegatus*

Matthew McShane  
*Illinois Wesleyan University*

William Jaeckle, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

McShane, Matthew and Jaeckle, Faculty Advisor, William, "Metabolic Contribution of Particulate Matter (<math>< \mu\text{m}</math>) for Larvae of *Lytechinus variegatus*" (2012). *John Wesley Powell Student Research Conference*. 12.

<https://digitalcommons.iwu.edu/jwprc/2012/posters2/12>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P24

**METABOLIC CONTRIBUTION OF PARTICULATE MATTER (<1 $\mu$ m) FOR LARVAE OF *LYTECHINUS VARIEGATUS***

Matthew McShane and William Jaeckle\*  
Biology Department, Illinois Wesleyan University

Echinoderm larvae possess an external band of cilia that beat in one direction for locomotion and undergo localized beat reversal when triggered by particles ( $\geq 2$  mm), pushing these particles towards the mouth for ingestion. Particles below adequate size, if ingested, must be collected by passing seawater through the digestive system. We measured the ability of pluteus larvae of *Lytechinus variegatus* to capture bacteria-sized particles after exposing them to 0.513 $\mu$ m fluorescent beads ( $9.5 \times 10^5$  beads/mL) for 10 min. Beads were found in the digestive system in 75% (41 of 55) of larvae and the average ingestion rate was 39.4 ( $\pm 42.7$  SD) beads/h. From these ingestion rates we calculated clearance rates (mL/ larva-h) and used literature values for bacterial energy content and larval metabolism to estimate the energetic contribution of bacteria-sized particles. Assuming 100% assimilation efficiency and  $10^6$  cells/mL, we found that feeding on bacteria-sized particles could provide for only 0.16% of metabolic rate.