



Apr 14th, 9:00 AM - 10:00 AM

## Scanning Electrochemical Microscopy Mapping of Neurotransmitter Release Sites from Model Neurons

Jennifer A. Schreiber  
*Illinois Wesleyan University*

Melinda Baur, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>

---

Schreiber, Jennifer A. and Baur, Faculty Advisor, Melinda, "Scanning Electrochemical Microscopy Mapping of Neurotransmitter Release Sites from Model Neurons" (2012). *John Wesley Powell Student Research Conference*. 17.  
<https://digitalcommons.iwu.edu/jwprc/2012/posters/17>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P33

**SCANNING ELECTROCHEMICAL MICROSCOPY MAPPING OF  
NEUROTRANSMITTER RELEASE SITES FROM MODEL NEURONS**

Jennifer A. Schreiber and Melinda Baur\*  
Chemistry Department, Illinois Wesleyan University

Scanning Electrochemical Microscopy (SECM) was used to image the topography of model neurons in the constant height mode while simultaneously measuring neurotransmitter concentration using amperometry in the collector mode. The probe of the SECM was a fabricated 5  $\mu\text{m}$  carbon-fiber electrode. The electrode was calibrated to determine the distance to the surface. Topographical imaging was accomplished by moving the electrode systematically across the surface. Multiple topographical images were obtained before and after stimulation of the model neurons with KCl. Neurotransmitters norepinephrine and dopamine were detected by setting the electrode to an oxidizing potential (+0.8 v). After stimulation, neurotransmitter release was detected amperometrically during topographical mapping. Download this file and use it as a template.