



Apr 16th, 11:00 AM - 12:00 PM

Effects of Exercise and Good Limb Training on Functional Outcome Following Stroke in C57bl/6 Mice

Amanda Macuiba
Illinois Wesleyan University

Abigail Kerr, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Education Commons](#), and the [Psychology Commons](#)

Macuiba, Amanda and Kerr, Faculty Advisor, Abigail, "Effects of Exercise and Good Limb Training on Functional Outcome Following Stroke in C57bl/6 Mice" (2016). *John Wesley Powell Student Research Conference*. 2.

<https://digitalcommons.iwu.edu/jwprc/2016/oralpres4/2>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation O4.2

**EFFECTS OF EXERCISE AND GOOD LIMB TRAINING ON
FUNCTIONAL OUTCOME FOLLOWING STROKE
IN C57BL/6 MICE**

Amanda Macuiba and Abigail Kerr*

Psychology Department, Illinois Wesleyan University

Stroke is a leading cause of long-term disability, making research on behavioral rehabilitation imperative. A common strategy following stroke is the compensatory use of the less impaired, or good, limb. Compensatory use of the good limb after stroke is efficient and promotes a quick return to independent daily living, but it ultimately negatively impacts overall recovery. We believe exercise may promote better recovery with compensatory limb use. Research shows that exercise promotes neuronal growth and prevents cell death. This study used mice to investigate whether exercise could prevent deterioration the bad limb associated with compensatory training of the good limb. Results showed that exercising mice, with or without good limb training, retained bad limb functionality and have better outcome than mice receiving compensatory training only. These findings suggest that exercise can maintain and extend the recovery potential of the bad limb while permitting compensatory limb use in the short term.