



Apr 16th, 11:00 AM - 12:00 PM

## Elucidating a Mechanism of Growth Cessation in Heterocysts of (*Anabaena*)

Blake Beehler  
*Illinois Wesleyan University*

Loralyn Cozy, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Biology Commons](#), and the [Education Commons](#)

---

Beehler, Blake and Cozy, Faculty Advisor, Loralyn, "Elucidating a Mechanism of Growth Cessation in Heterocysts of (*Anabaena*)" (2016). *John Wesley Powell Student Research Conference*. 3.

<https://digitalcommons.iwu.edu/jwprc/2016/oralpres6/3>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Oral Presentation O6.3

**ELUCIDATING A MECHANISM OF GROWTH CESSATION  
IN HETEROCYSTS OF *ANABAENA***

Blake Beehler and Loralyn Cozy\*  
Biology Department, Illinois Wesleyan University

*Anabaena* sp. strain PCC 7120 is a filamentous, multicellular cyanobacterium that serves as a model for the study of terminal cell type differentiation. When starved for bioavailable nitrogen, a subset of cells in a filament terminally differentiate into a nitrogen-fixing cell type, called a heterocyst. To maintain their position in the growing filament, heterocysts lose the ability to divide. This cell-type specific cessation of division suggests that there is a regulatory link between binary fission and heterocyst differentiation. It is unknown how cell division is inhibited in heterocysts, but it has been shown that heterocysts lack polymerized FtsZ rings (Z-rings) normally responsible for cytokinesis. Here, we find that a strain of *Anabaena* mutant for the *hetP* gene exhibited Z rings in heterocysts. This indicates that the *hetP* gene may directly or indirectly inhibit Z ring formation in heterocysts, thus preventing division. To test these two possibilities, the interactions between HetP and FtsZ were qualitatively and quantitatively assessed through a bacterial two-hybrid system and  $\beta$ -galactosidase assay, and a potential model relating the action of HetP and FtsZ was developed.