



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2016, 27th Annual JWP Conference

Apr 16th, 9:00 AM - 10:00 AM

Anatomical Effects of Exercise Following Ischemic Insult in Young and Aged C57BL/6 Mice

Mark Curtis
Illinois Wesleyan University

Abigail Kerr, Faculty Advisor
Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Education Commons](#), and the [Psychology Commons](#)

Curtis, Mark and Kerr, Faculty Advisor, Abigail, "Anatomical Effects of Exercise Following Ischemic Insult in Young and Aged C57BL/6 Mice" (2016). *John Wesley Powell Student Research Conference*. 4.

<https://digitalcommons.iwu.edu/jwprc/2016/posters/4>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P7

**ANATOMICAL EFFECTS OF EXERCISE FOLLOWING
ISCHEMIC INSULT IN YOUNG AND AGED C57BL/6 MICE**

Mark Curtis and Abigail Kerr*

Psychology Department, Illinois Wesleyan University

Stroke is a leading cause of long-term disability. Current rehabilitative strategies are expensive and often fail to yield complete recovery. Focused training of the impaired limb improves outcome in rodents, but these strategies require intensive training that is not feasible for humans. Because aerobic exercise has been found to induce beneficial changes in the brain, it is a promising rehabilitative strategy following stroke. Exercise may require less intensity and is less expensive than traditional therapy. The current study investigated the effect of post-stroke exercise on young and aged mice. Mice were trained on a skilled reaching task before receiving a focal ischemic stroke. Mice were subdivided into three different groups for rehabilitative training: traditional rehabilitation, aerobic exercise, and control procedures. Both young and aged mice benefited from aerobic exercise after stroke. Aerobic exercise may be an affordable and effective alternative to traditional rehabilitative strategies. Underlying anatomical mechanisms are currently being analyzed.