



Apr 16th, 2:00 PM - 3:00 PM

# Functional Complementation of Z-Ring Regulation by Alleles of *HETP* in *Anabaena*

Christina Khour

*Illinois Wesleyan University*

Loralyn Cozy, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

 Part of the [Biology Commons](#), and the [Education Commons](#)

Khour, Christina and Cozy, Faculty Advisor, Loralyn, "Functional Complementation of Z-Ring Regulation by Alleles of *HETP* in *Anabaena*" (2016). *John Wesley Powell Student Research Conference*. 9.  
<http://digitalcommons.iwu.edu/jwprc/2016/posters2/9>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P18

## FUNCTIONAL COMPLEMENTATION OF Z-RING REGULATION BY ALLELES OF *HETP* IN *ANABAENA*

Christina Khouri and Loralyn Cozy\*  
Biology Department, Illinois Wesleyan University

*Anabaena* is a filamentous cyanobacterium that is able to undergo cell-differentiation in nitrogen-limiting conditions to produce a nitrogen-fixing cell called a heterocyst. Under normal conditions, *Anabaena* will produce 10% heterocysts in its filament in about 23 hours. While the vegetative cells continue to divide by constriction of a protein ring, called the Z-ring, heterocysts do not contain z-rings and do not divide. In contrast, a  $\Delta$ *hetP* mutant of *Anabaena* produces only 2-3% heterocysts at a later time of 48 hours, with z-rings present in about one-third of these heterocysts. The part of the *hetP* gene required for inhibition of z-ring formation in heterocysts is unknown. To identify this part of the gene, we attempted to complement  $\Delta$ *hetP* mutant by the addition alleles of the *hetP* gene. Three alleles of *hetP* were tested in comparison to a  $\Delta$ *hetP* mutant containing empty vector alone: a wild-type copy of *hetP*, *hetP* containing the amino acid mutations C36A and C95A, and a 68 base pair truncation of *hetP*. The presence of Z-rings in heterocysts was assessed after 48 hours in nitrogen-limiting conditions by fluorescence microscopy. Z-rings in heterocysts were only found in the  $\Delta$ *hetP* strain of *Anabaena* containing an empty vector suggesting a limited region of the *hetP* gene that is required for z-ring loss in heterocysts.

### References:

1. Higa, K. C., Callahan, S. M.. (2010) Ectopic expression of *hetP* can partially bypass the need for *hetR* in heterocyst differentiation by *Anabaena* sp. Strain PCC 7120. *Molecular Microbiology* 77(3): 564-574.

