



Illinois Wesleyan University
Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2016, 27th Annual JWP Conference

Apr 16th, 9:00 AM - 10:00 AM

c7 and C7 Complement Multidecomposition of Kn

Hang Phung

Illinois Wesleyan University

Daniel Roberts, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Education Commons](#), and the [Mathematics Commons](#)

Phung, Hang and Roberts, Faculty Advisor, Daniel, "c7 and C7 Complement Multidecomposition of Kn" (2016). *John Wesley Powell Student Research Conference*. 14. <https://digitalcommons.iwu.edu/jwprc/2016/posters/14>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P31

C_7 AND C_7 COMPLEMENT MULTIDECOMPOSITION OF K_n

Hang Phung and Daniel Roberts*

Mathematics Department, Illinois Wesleyan University

If all edges of K_n can be partitioned into copies of a graph G , such design is called a G -decomposition of K_n , or G -design of order n . The extension of this design, called multi-decomposition, is decomposition of K_n by a graph pair. For any integer $v \geq 4$, a graph-pair of order v is a pair of non-isomorphic graphs G and H of order v such that there are no isolated vertices and $E(G) \cup E(H) = E(K_v)$. A $(G; H)$ –multi-decomposition of K_n is determined when all edges of K_n can be partitioned into copies of G and H with at least a copy of either G or H . In the past, the multi-designs for all graph-pairs of order 4 and 5 have been finished and published in 2003. The necessary and sufficient conditions for the existence of a $(C_6; \overline{C_6})$ – multi-decomposition of K_n has also been found out by Gao Yizhe. This paper is dedicated to continuing the project by determining the condition for n such that there exists $(C_7, \overline{C_7})$ –multi-decomposition of K_n .

Reference

Abueida, A. A., and Daven, M. Multidesigns for graph-pairs of order 4 and 5. *Graphs and Combinatorics*, 19(4), (2003) 433-447.