



**Illinois Wesleyan University**  
**Digital Commons @ IWU**

---

John Wesley Powell Student Research  
Conference

2016, 27th Annual JWP Conference

---

Apr 16th, 9:00 AM - 10:00 AM

## Restoration of Heterocyst Production to a $\Delta$ HETP Strain of *Anabaena*

Megan Smeets  
*Illinois Wesleyan University*

Loralyn Cozy, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <https://digitalcommons.iwu.edu/jwprc>



Part of the [Biology Commons](#), and the [Education Commons](#)

---

Smeets, Megan and Cozy, Faculty Advisor, Loralyn, "Restoration of Heterocyst Production to a  $\Delta$ HETP Strain of *Anabaena*" (2016). *John Wesley Powell Student Research Conference*. 15.

<https://digitalcommons.iwu.edu/jwprc/2016/posters/15>

This Event is protected by copyright and/or related rights. It has been brought to you by Digital Commons @ IWU with permission from the rights-holder(s). You are free to use this material in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself. This material has been accepted for inclusion by faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P33

## **RESTORATION OF HETEROCYST PRODUCTION TO A $\Delta$ HETP STRAIN OF ANABAENA**

Megan Smeets and Loralyn Cozy\*

Biology Department, Illinois Wesleyan University

*Anabaena* sp. Strain PCC 7120 is a filamentous cyanobacterium capable of differentiating a nitrogen-fixating cell type called a heterocyst. The *hetP* gene has been identified as being required for the normal magnitude and timing of heterocyst formation. Where as a wild type strain produces 10% heterocysts in 23 hours, a  $\Delta$ *hetP* strains produces only 2-3% heterocysts in 48 hours. How the loss of *hetP* leads to this phenotype is currently unknown. To identify genes downstream of *hetP* in the differentiation pathway that could be aiding in its function during heterocyst formation we performed a forward genetic screen. A Tn5 transposon was introduced via conjugation into a  $\Delta$ *hetP* strain and plated on nitrogen deficient media with selection. Surviving colonies were assayed for restoration of wild-type heterocyst accumulation and timing of development. The DNA of these strains was isolated and the transposon location was identified. We demonstrated a class of genes whose inactivation is capable of restoring wild-type heterocyst formation in the absence of *hetP*. Future work will examine the function of these genes in relation to *hetP*.