



May 8th, 9:30 AM - 4:30 PM

The Effects of the Hematophagous Mite (*Dermanyssus hirundinis*) on Nestling House Wrens (*Troglodytes aedon*)

Jennifer L. Hoger

Illinois Wesleyan University

Andrew Pacejka

Illinois State University

Given Harper, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Hoger, Jennifer L.; Pacejka, Andrew; and Harper, Faculty Advisor, Given, "The Effects of the Hematophagous Mite (*Dermanyssus hirundinis*) on Nestling House Wrens (*Troglodytes aedon*)" (1993). *John Wesley Powell Student Research Conference*. 36.

<http://digitalcommons.iwu.edu/jwprc/1993/posters/36>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

THE EFFECTS OF THE HEMATOPHAGOUS MITE (*Dermanyssus hirundinis*)
ON NESTLING HOUSE WRENS (*Troglodytes aedon*).

Jennifer L. Hoger, Biology Dept., IWU, Andrew Pacejka, ISU, R. Given Harper*.

The house wren (*Troglodytes aedon*) is a small, insectivorous, migratory passerine bird that nests in secondary tree cavities and nestboxes. Both hematophagous (blood-sucking) mites, *Dermanyssus hirundinis*, and predatory (non-blood-sucking) mites, *Androlalaps casalis*, have been observed in the nesting material of the house wren. However, no studies have investigated their effects on nestling house wrens. In other avian species such ectoparasites have had adverse effects on nestlings, including low nestling mass, premature fledging (i.e. nest leaving) dates, and low nestling survival.

The purpose of my study was to determine the effects of *Dermanyssus hirundinis* on the mass at fledging and survival of nestling house wrens. The study was conducted from 1990-1992 in a population of house wrens that bred in nestboxes (N=910) near Bloomington, IL. Mites were extracted from the nesting material in the nestboxes from May-August of each year, and were then counted. These data were then correlated to nestling mass and the date of fledging. The results of this study will be presented at the research conference.