



May 8th, 9:30 AM - 4:30 PM

# An Animals Model of Alzheimer's Disease III: Characterization of Neuronal Degeneration Using the Technique of Immunocytochemistry for Glial Fibrillary Acidic Protein

Esther Kang

*Illinois Wesleyan University*

Wayne Dornan, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Kang, Esther and Dornan, Faculty Advisor, Wayne, "An Animals Model of Alzheimer's Disease III: Characterization of Neuronal Degeneration Using the Technique of Immunocytochemistry for Glial Fibrillary Acidic Protein" (1993). *John Wesley Powell Student Research Conference*. 30.  
<http://digitalcommons.iwu.edu/jwprc/1993/posters/30>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

AN ANIMALS MODEL OF ALZHEIMER'S DISEASE III:  
CHARACTERIZATION OF NEURONAL DEGENERATION USING THE  
TECHNIQUE OF IMMUNOCYTOCHEMISTRY FOR  
GLIAL FIBRILLARY ACIDIC PROTEIN

Esther Kang, Dept of Natural Science, IWU, Dr. Wayne A. Dornan

Alzheimer's disease (AD) currently affects an estimated 4 million Americans. Manifested initially by mild forgetfulness, this devastating disease eventually erodes all cognitive abilities. Neuropathologically AD is characterized by neuritic plaques (NP) and intracellular neurofibrillary tangles (NFT). Although the etiology of AD is unknown, NP deposition has been suggested to play a fundamental role. Recently, the major component of NP has been identified as a B-amyloid protein that exists in an insoluble state. This pathology is invariably accompanied by the proliferation of adjacent glial cells in response to neuronal degeneration. In particular, postmortem studies of AD patients show a marked increase in the level of glial fibrillary acidic protein (GFAP) due to astrogliosis. Moreover, recent development of immunocytochemical techniques which stain specific proteins, have shown strong correlation between the degree of dementia in AD patients with the degree of intensity of GFAP staining. These results strongly suggest that immunocytochemistry for GFAP is a reliable tool in measuring neurodegeneration in AD patients.

In a study that is being conducted concurrently (see David Kang's abstract), intrahippocampal injections of Beta-Amyloid (25-35) or Ibotenic Acid have been used to induce a potential animal model of AD. In this study GFAP immunocytochemistry will be employed to assess the degree of neuronal degeneration in these animals. The result of this assessment will be presented at the conference.