



Apr 15th, 2:00 PM - 3:00 PM

Applying Van Der Pauw's Technique to Thermal Conductivity

Mark Ordonez

Illinois Wesleyan University

Ileana Rau

Illinois Wesleyan University

Gabriel C. Spaulding, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Ordonez, Mark; Rau, Ileana; and Spaulding, Faculty Advisor, Gabriel C., "Applying Van Der Pauw's Technique to Thermal Conductivity" (2000). *John Wesley Powell Student Research Conference*. 14.

<http://digitalcommons.iwu.edu/jwprc/2000/posters2/14>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation 36

**APPLYING VAN DER PAUW'S TECHNIQUE TO
THERMAL CONDUCTIVITY**

Mark Ordonez and Ileana Rau and Gabe Spalding*
Department of Physics, Illinois Wesleyan University

When measuring electrical resistance, two leads (an input and an output) may be used so long as the resistance of the sample is large compared to that of the leads and the contacts, and as long as thermoelectric voltages are negligible in comparison to the signal of interest. However, multi-lead configurations can often provide further information. For example, a method due to van der Pauw can be used to extract the intrinsic electrical resistivity of a material even from samples having non-trivial geometry (where current flow is non-homogeneous).

Formally, there is a great similarity between electronic and heat transport. In 1999 researchers at the University of Freiburg, Germany took advantage of this similarity, and applied the van der Pauw technique to measurements of thermal conductivity. Because such a method provides a more detailed mapping of the thermal conductivity, we intend to explore the use of the van der Pauw technique to map out the intrinsic anisotropies in the thermal conductivity of a single crystal sample.

Moreover, by exploring the thermal conductivity of SrTiO_3 samples at very low temperatures (down to ~ 0.5 K), it may be possible to help explain some of the peculiar quantum effects seen in this material.