



Apr 8th, 9:00 AM - 10:00 AM

## Organochlorine Pesticide Contamination in Resident North American Passerines

Amy Cadwallader

*Illinois Wesleyan University*

Kevin Latman,

*Illinois Wesleyan University*

Given Harper, Faculty Advisor

*Illinois Wesleyan University*

Jeffrey Frick, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Cadwallader, Amy; Latman,, Kevin; Harper, Faculty Advisor, Given; and Frick, Faculty Advisor, Jeffrey, "Organochlorine Pesticide Contamination in Resident North American Passerines" (2006). *John Wesley Powell Student Research Conference*. 6.

<http://digitalcommons.iwu.edu/jwprc/2006/posters/6>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P15

**ORGANOCHLORINE PESTICIDE CONTAMINATION IN RESIDENT  
NORTH AMERICAN PASSERINES**

Amy Cadwallader and Kevin Latman and Given Harper and Jeffrey Frick\*  
Biology and Chemistry Departments, Illinois Wesleyan University

Organochlorine (OC) pesticide and metabolite levels were determined in resident North American passerines and woodpeckers collected in McLean County, Illinois. Sixty-nine of 77 individual birds contained OC compounds above detection limits, including all eighteen species examined. Total contamination levels in individual birds ranged from 7.47 to 2,274.23 ng/g. The most prevalent OC compound was p,p'-DDE, which was present in 45 birds; dieldrin was found in 30 birds, heptachlor and heptachlor epoxide were each found in 27 birds, and p,p'-DDT was found in 25 birds. There was no significant difference in total OC levels between males and females, between HY and AHY birds, and between birds that bred and wintered in more northern latitudes. There was a significant difference in total OC levels between diet categories. Omnivores had significantly higher total OC levels than granivores. House sparrows had significantly lower total OC levels than both white-throated sparrows and American robins.