



Apr 9th, 9:00 AM - 10:00 AM

# A New Species of *Pristimantis* (Amphibia: Anura: Strabomantidae) from the Río Abiseo National Park, Peru

Cindy Gregory

*Illinois Wesleyan University*

Edgar Lehr, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

 Part of the [Biology Commons](#)

Gregory, Cindy and Lehr, Faculty Advisor, Edgar, "A New Species of *Pristimantis* (Amphibia: Anura: Strabomantidae) from the Río Abiseo National Park, Peru" (2011). *John Wesley Powell Student Research Conference*. 9. <http://digitalcommons.iwu.edu/jwprc/2011/posters/9>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P17

**A NEW SPECIES OF *PRISTIMANTIS* (AMPHIBIA: ANURA: STRABOMANTIDAE)  
FROM THE RÍO ABISEO NATIONAL PARK, PERU**

Cindy Gregory and Edgar Lehr\*  
Biology Department, Illinois Wesleyan University

Frogs of the genus *Pristimantis* occur in Central America, the Lesser Antilles, and northwestern South America. They reproduce by terrestrial-eggs without a free-swimming tadpole stage. *Pristimantis* are most diverse in South America and can be found from lowland forests to elevations of about 4000 m in the Andes. The genus comprises 440 species, of which 120 are currently known from Peru. Herein, we describe a new species of *Pristimantis* from the Río Abiseo National Park in the Andes of northern Peru. The new species has a snout-vent length of 24.9–34.2 mm (n = 7) in adult females, and 15.3–23.5 mm (n = 32) in adult males. It differs from other species of *Pristimantis* in having the snout with a broad, slightly upwards curved, fleshy process. It is most similar to *P. phoxocephalus* which has the snout with a vertical fleshy keel, but differs from it by a combination of morphological characters.